

September 1999

Finite Temperature Time-Dependent Effective Theory For The Goldstone Field In A BCS-Type Superfluid

Ian J.R. Aitchison ¹, Georgios Metikas ²

Department of Physics, Theoretical Physics,
University of Oxford, 1 Keble Road, Oxford OX1 3NP

and

Dominic J. Lee ³

Department of Physics,
University of Leeds, Leeds LS2 9JT

Abstract

We extend to finite temperature the time-dependent effective theory for the Goldstone field (the phase of the pair field) θ which is appropriate for a superfluid containing one species of fermions with s-wave interactions, described by the BCS Lagrangian. We show that, when Landau damping is neglected, the effective theory can be written as a local time-dependent non-linear Schrödinger Lagrangian (TDNLSL) which preserves the Galilean invariance of the zero temperature effective theory and is identified with the superfluid component. We then calculate the relevant Landau terms which are non-local and which destroy the Galilean invariance. We show that the retarded θ -propagator (in momentum space) can be well represented by two poles in the lower-half frequency plane, describing damping with a predicted temperature, frequency and momentum dependence. It is argued that the real parts of the Landau terms can be approximately interpreted as contributing to the normal fluid component.

¹E-mail address: i.aitchison1@physics.oxford.ac.uk

²E-mail address: g.metikas1@physics.oxford.ac.uk

³E-mail address: phydjl@phys-irc.novell.leeds.ac.uk

1 Introduction

The form of the effective (Ginzburg-Landau) Lagrangian for time-dependent superconducting phenomena continues to attract attention. At $T = 0$ Schakel [1], using derivative expansion techniques [2], obtained a time-dependent effective Lagrangian for the Goldstone field (which is the phase θ of the pair field Δ), starting from a simple BCS Lagrangian. Interestingly, Schakel's effective Lagrangian is the same as one that had been proposed earlier [3] by assuming that the momentum density in a Galilean invariant system is proportional to a conserved current. In [4] (see also [5]) the Lagrangian of [1] and [3] was shown to be equivalent, up to a certain order in derivatives, to a time-dependent non-linear Schrödinger Lagrangian (TDNLSL); in particular, the role of Galilean invariance was emphasized. More recently, de Palo et al. [6] have given a more general derivation of the TDNLSL at $T = 0$, by including in the effective action from the outset fluctuations in the density field ρ (conjugate to θ), as well as those in Δ . This formulation allows a unified description of the weak-coupling (BCS) and the strong-coupling (Bose) limits as well as of the cross-over between them [6].

In all the above cases, at $T = 0$, electromagnetic interactions can be straightforwardly included by the “minimal coupling” prescription, since the effective Lagrangian is local. As is well-known [7], the $q^2 = 0$ pole of the θ -mode is then pushed to finite frequencies, becoming the plasma mode.

At finite temperature, in the non-static case, the situation is complicated by the presence of Landau damping terms, which appear to prevent *any* local time-dependent Lagrangian for θ , since they have singularities at $q_0 = \pm v_F |\mathbf{q}|$ (see Section 6 below), and the resulting cut structure implies that there is no Taylor series expansion about the origin in momentum space. This problem was first pointed out by Abrahams and Tsuneto [8]. It poses a fundamental difficulty for the inclusion of electromagnetism, since minimal coupling cannot be used unless the effective Lagrangian is local. Note that the singularities, and hence this difficulty, disappear in the static case, so that there is, of course, nothing wrong with the familiar static Ginzburg-Landau theory at $T \neq 0$.

Adopting a modern field-theoretical approach to the problem of deriving an effective Lagrangian, Stoof [9] neglected the Landau damping terms entirely and constructed a time-

dependent effective theory by writing the pair field as $\Delta(\mathbf{x}, t) = |\Delta_0| + \Delta'(\mathbf{x}, t)$, where $|\Delta_0|$ is the value of Δ which minimizes the effective action, and $\Delta'/|\Delta_0|$ was assumed to be varying slowly in space and time. In [10] an attempt was made to extend Stoof's approach so as to include Landau damping, retaining only terms quadratic in Δ' . It was argued that the troublesome (Landau) cut structure could be replaced - in the θ -propagator - by two simple poles at complex energy. This led to an approximate *local* effective action for θ of the form

$$\int d^4x \left[\dot{\theta}^2 - [a(T) - ib(T)]v_F^2(\nabla\theta)^2 \right] \quad (1)$$

where $a(T)$, $b(T)$ are real functions. If (1) were correct, minimal coupling could, as usual, be employed to include electromagnetism.

Unfortunately (1) is *not* correct. The imaginary part of the Goldstone mode propagator, S_G , is an odd function of the frequency and will not contribute to the total effective action as in (1) (it does, however, enter the equation of motion for θ). The correct approximate two-pole form of S_G will be given in Section 7 below, and it will be such that S_G^{-1} is not local in coordinate space. Thus we are back with the problem of how to include electromagnetism in the non-static case for $T \neq 0$.

Since we do not at present know of any answer to this problem, all our subsequent analysis will generally refer to BCS-type pairing of *neutral* fermions, i.e. to an uncharged fermionic superfluid. Helium-3 is such a fluid but it condenses into a p-wave state, not the s-wave BCS state studied here. It would be straightforward, and interesting, to generalize the present approach to p-wave pairing (a beginning was made by Van Weert [11]). However, it is likely that trapped condensates of neutral fermionic fluids will be produced before very long (see for example [12]). Our analysis would be relevant to such a case, when re-done for a finite system.

Apart from the error in S_G , the approach leading to (1) is unsatisfactory for another reason, namely it fails to reduce, at $T = 0$, to the correct Galilean-invariant result of [1, 3, 4]. The problem can be traced to the step $\Delta = |\Delta_0| + \Delta'$: as was pointed out in [4], this ansatz destroys Galilean invariance and fails to allow for large fluctuations in θ , two weaknesses which are in fact related. In [10], the further approximations $\Delta \approx |\Delta| (1 + i\theta)$ and $|\Delta| = \text{constant}$ were made, which led to the form (1) after retaining only quadratic terms in $\text{Im}(\Delta)$. What is required, rather, is the finite temperature version of [4] (or even better of [6]), where we write

$\Delta = |\Delta| e^{i\theta}$, ignore fluctuations in $|\Delta|$ and consider a systematic expansion in the derivatives of θ , without having to assume that θ itself is small.

Our aim in this paper is to provide the finite temperature extension of [4] and arrive at a temperature and time-dependent effective Lagrangian for θ , $L(\theta)$. In Section 2 we set up the effective action formalism and derive the usual gap equation. In Section 3 we evaluate the first term in the expansion of the effective action in powers of derivatives and in Section 4 we calculate the next term which includes the Landau damping contribution. In Section 5 we show that, if the Landau terms are neglected, Galilean invariance is maintained (at least up to the order in derivatives to which we are working) and $L(\theta)$ has exactly the same form as in [1,3,4] but with temperature dependent coefficients. This, in turn, implies that an effective non-linear Schrödinger theory can be obtained with a temperature-dependent potential term.

Next, we turn our attention to the Landau terms, which are calculated in momentum space in Section 6. As expected, their form is similar to the expressions found in [10]. Their presence prevents any representation of the effective action in terms of a local Lagrangian. In Section 7 we consider how to obtain a useful approximate form for the equation of motion for θ , including the Landau terms. Noting (see above) that $\text{Im}(S_G)$ is an odd function of frequency, we show that S_G can be well represented by two poles in the lower half plane of the complex frequency variable. This corrects the form proposed in [10] and invalidates (1). We then consider the real parts of the Landau terms, neglecting the imaginary parts. We propose that they can be approximated by their value at the (undamped) poles of S_G - an “on-shell” approximation. The complete effective action then becomes local, though the Landau terms break Galilean invariance. We propose a physical interpretation of the resulting equation of motion, in which the real parts of the Landau terms represent a contribution to the normal fluid component.

2 The Effective Action and the Gap Equation

We begin with a brief review of the formalism. The BCS Lagrangian, for s-wave pairing and in the absence of external fields, is

$$L = \sum_{\sigma=1}^2 \psi_{\sigma}^*(x) \left(i\partial_t + \frac{\nabla^2}{2m} + \mu \right) \psi_{\sigma}(x) + g \psi_{\uparrow}^*(x) \psi_{\downarrow}^*(x) \psi_{\downarrow}(x) \psi_{\uparrow}(x) \quad (2)$$

where $\psi_{\sigma}(x)$ describes electrons with spin $\sigma = (\uparrow, \downarrow)$, $\mu = \frac{k_F^2}{2m}$ is the Fermi energy, $x = (t, \mathbf{x})$ and $g > 0$. We define the Nambu spinor

$$\psi = \begin{pmatrix} \psi_{\downarrow} \\ \psi_{\uparrow}^* \end{pmatrix}$$

and introduce the auxiliary “pair” fields $\Delta(x)$ and $\Delta^*(x)$ so as to write the partition function as

$$Z = \int D\psi^{\dagger} D\psi \int D\Delta^* D\Delta \exp \left[i \int d^4x \left(\psi^{\dagger} \mathcal{G}_{xx}^{-1} \psi - \frac{|\Delta|^2}{g} \right) \right] \quad (3)$$

where

$$\mathcal{G}_{xx}^{-1} = \begin{pmatrix} i\partial_t + \frac{\nabla^2}{2m} + \mu & \Delta \\ \Delta^* & i\partial_t - \frac{\nabla^2}{2m} - \mu \end{pmatrix}.$$

We now do a gauge transformation on the electron field [1], writing $\psi_{\sigma} = \exp\left(\frac{i\theta(x)}{2}\right) \chi_{\sigma}(x)$, where θ is the phase of Δ and χ_{σ} are new anticommuting fields. The integrals over χ and $\chi^{\dagger}$ are then performed, leading to

$$Z = \int D\Delta^* D\Delta \exp(iS_{eff}[\Delta, \Delta^{\dagger}]) \quad (4)$$

where the effective action S_{eff} is given by

$$S_{eff}[\Delta, \Delta^*] = -i\text{Tr} \log(\hat{G}^{-1} - \hat{\Sigma}) - \int d^4x \frac{|\Delta|^2}{g} \quad (5)$$

$$= -i\text{Tr} \log \hat{G}^{-1} + i\text{Tr} \sum_{n=1}^{\infty} \frac{(\hat{G}\hat{\Sigma})^n}{n} - \int d^4x \frac{|\Delta|^2}{g} \quad (6)$$

with

$$\begin{aligned}
\hat{G}^{-1} &= \begin{pmatrix} \hat{p}_0 - \frac{\hat{\mathbf{p}}^2}{2m} + \mu & |\hat{\Delta}| \\ |\hat{\Delta}| & \hat{p}_0 + \frac{\hat{\mathbf{p}}^2}{2m} - \mu \end{pmatrix} \\
\hat{\Sigma} &= \left(-\frac{i\widehat{\nabla^2\theta}}{4m} + \frac{(\widehat{\nabla\theta})\hat{\mathbf{p}}}{2m} \right) I + \left(\frac{\hat{\theta}}{2} + \frac{(\widehat{\nabla\theta})^2}{8m} \right) \tau_3 \\
\hat{f} &= f(\hat{x}).
\end{aligned}$$

Note that the effect of the gauge transformation has been to separate the dependence on $|\hat{\Delta}|$ which is in $\hat{G}^{-1}$ from that on $\hat{\theta}$ which is in $\hat{\Sigma}$. We point out that Det and Tr refer to both the functional space and the internal (Nambu) space. The operators $\hat{p}_0$, $\hat{\mathbf{p}}$ and $\hat{x}$ are Fock-Schwinger Proper Time operators.

The ground state is determined by taking $|\Delta|$ to be space-time independent and solving the mean field equation

$$\frac{\delta S_{eff}}{\delta |\Delta|} = 0 \Leftrightarrow ig \int \frac{d^4 p}{(2\pi)^4} \frac{1}{p_0^2 - E^2} = 1 \quad (7)$$

where $E^2 = \epsilon^2 + |\Delta|^2$ and $\epsilon = \frac{\mathbf{p}^2}{2m} - \mu$. The modulus $|\Delta|$ of the pair field is therefore the *energy gap*. (7) can be written in the usual form of the BCS gap equation at finite temperature [7, 13]

$$g \int \frac{d^3 \mathbf{p}}{(2\pi)^3} \frac{1}{2E} \tanh \frac{\beta E}{2} = 1 \quad (8)$$

where we assume that the potential (the coupling constant) is attractive in a shell of width $2\omega_c$ centered at the Fermi surface and vanishes outside this shell. We note that in the weak-coupling limit $\omega_c \gg |\Delta(0)|$, where $|\Delta(0)|$ refers to the value of the gap within the shell. In this limit we can consider $|\Delta|$ to be a time and space independent constant whose value is

$$|\Delta(T)| \simeq |\Delta(0)| \simeq 1.75 k_B T_c \quad (9)$$

for

$$b = T/T_c \in [0.1, 0.6]. \quad (10)$$

Thus the effective theory only involves the single degree of freedom θ . If we expand each term of the infinite series in (6) in powers of momentum and frequency and restrict ourselves to

small values of these quantities, it has been proved in [1, 4] that at $T = 0$ the trace of the first term contributes a term proportional to $[\dot{\theta} + (\nabla\theta)^2/4m]$ to $L(\theta)$, and the trace of the second term contributes a term proportional to $[\dot{\theta} + (\nabla\theta)^2/4m]^2$. As emphasised in [4], the combination $[\dot{\theta} + (\nabla\theta)^2/4m]$ is Galilean invariant and it is this feature which allows $L(\theta)$ to be rewritten as a TDNLSL at zero temperature. We shall now extend this calculation to finite temperature and explore to what extent an effective Lagrangian is still derivable and - if so - whether it can be written as a TDNLSL.

3 The First Trace

In this section we calculate the first ($n = 1$) term of the sum appearing in (6) :

$$\text{Tr}[\hat{G}\hat{\Sigma}] = \int \frac{d^4p}{(2\pi)^4} \text{tr}[G(p_0, \mathbf{p})\tau_3] \int d^4x \frac{1}{2} [\dot{\theta} + \frac{(\nabla\theta)^2}{4m}] \quad (11)$$

where a term odd in p_0 has vanished in the last step, since we integrate over p_0 from $-\infty$ to $+\infty$. By tr we mean the trace only in the internal (Nambu) space. We apply the Matsubara formalism to evaluate the first integral at finite temperature. We Wick-rotate via $p_0 \rightarrow ip_{0E}$ and then replace the integral over p_{0E} by a sum over discrete fermion frequencies via

$$i \int_{-\infty}^{\infty} \frac{dp_{0E}}{2\pi} f(ip_{0E}) \rightarrow \frac{i}{\beta} \sum_{n=-\infty}^{\infty} f(iz_n) \quad (12)$$

where $z_n = (2n + 1)\pi/\beta$, and n is integer. Next, following [14], we write

$$\frac{i}{\beta} \sum_n f(iz_n) \equiv \frac{i}{\beta} \sum_n f_E(z_n) = \frac{-1}{4\pi} \int_{C_R} dz f_E(z) \tan\left(\frac{\beta z}{2}\right) \quad (13)$$

where the contour C_R encloses the real axis in a counter-clockwise sense. (13) is valid provided that $f_E(z)$ has no singularities on the real axis, which is our case. It is convenient to rewrite the above equation in such a way as to separate out the (known) $T = 0$ contribution :

$$\frac{-1}{4\pi} \int_{C_R} dz f_E(z) \tan\left(\frac{\beta z}{2}\right) = L1 + L2 + L3 \quad (14)$$

with

$$\begin{aligned}
L1 &= \frac{i}{2\pi} \int_{-\infty}^{\infty} dz f_E(z) \\
L2 &= -\frac{i}{2\pi} \int_{-\infty+i\eta}^{\infty+i\eta} dz f_E(z) n(-iz) \\
L3 &= -\frac{i}{2\pi} \int_{-\infty-i\eta}^{\infty-i\eta} dz f_E(z) n(iz)
\end{aligned} \tag{15}$$

where η is an infinitesimal positive quantity and $n(x) = \frac{1}{1+\exp \beta x}$ is the Fermi-Dirac distribution which has been introduced in our calculation through the use of the identities $\tan(\frac{\beta z}{2}) = -i[1 - 2n(iz)] = i[1 - 2n(-iz)]$.

Applying the above formalism, we find that the total contribution of the first term of the series for S_{eff} is

$$i\text{Tr}[\hat{G}\hat{\Sigma}] = - \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[1 - \frac{\epsilon}{E} [1 - 2n(E)] \right] \int d^4x \frac{1}{2} [\dot{\theta} + \frac{(\nabla\theta)^2}{4m}] \tag{16}$$

which has the same Galilean-invariant form as at $T = 0$ but with a temperature dependent coefficient. In fact, the temperature dependence of the first trace is not important and can be neglected. The maximum value of the function $n(E)$ for the range of temperatures of (10) and with the gap having the value given in (9) is $n_{max} \simeq 0.05$. Consequently $1 - 2n(E) \simeq 1$ and the coefficient of the Galilean invariant term $[\dot{\theta} + \frac{(\nabla\theta)^2}{4m}]$ is

$$M(T) = \frac{1}{2} \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[1 - \frac{\epsilon}{E} [1 - 2n(E)] \right] \simeq \frac{1}{2} \rho_0 = M(0), \tag{17}$$

where $\rho_0 = p_F^3/(3\pi^2)$ is the electron density at $T = 0$.

4 The Second Trace

The trace of the second term ($n = 2$) of the infinite series in (6) can be written as:

$$\text{Tr}[\hat{\Sigma}\hat{G}\hat{\Sigma}\hat{G}] = \text{tr} \int \frac{d^4p}{(2\pi)^4} \frac{d^4q}{(2\pi)^4} \tilde{\Sigma}(q, \mathbf{p} - \frac{\mathbf{q}}{2}) G(p_0 - \frac{q_0}{2}, \mathbf{p} - \frac{\mathbf{q}}{2}) \tilde{\Sigma}(-q, \mathbf{p} + \frac{\mathbf{q}}{2}) G(p_0 + \frac{q_0}{2}, \mathbf{p} + \frac{\mathbf{q}}{2}) \tag{18}$$

where $\tilde{\Sigma}(p, \mathbf{q}) = \int d^4x \exp(ip.x) \Sigma(x, \mathbf{q})$. After performing the Wick rotation we have

$$\text{Tr}[\hat{\Sigma}\hat{G}\hat{\Sigma}\hat{G}]_E = - \int \frac{d^4q_E}{(2\pi)^4} \frac{d^4p_E}{(2\pi)^4} T_1 - 2i \int \frac{d^4q_E}{(2\pi)^4} \frac{d^4k_E}{(2\pi)^4} \frac{d^4p_E}{(2\pi)^4} T_2 + \int \frac{d^4q_E}{(2\pi)^4} \frac{d^4k_E}{(2\pi)^4} \frac{d^4r_E}{(2\pi)^4} \frac{d^4p_E}{(2\pi)^4} T_3 \tag{19}$$

where

$$\begin{aligned}
T_1 &= \tilde{\theta}_E(q_E) \tilde{\theta}_E(-q_E) A_{E-} A_{E+} \left[B_E \left[\frac{(\mathbf{p} \cdot \mathbf{q})^2}{2m^2} + \frac{q_{0E}^2}{2} \right] + C_E \left[-\frac{(\mathbf{p} \cdot \mathbf{q})^2}{2m^2} + \frac{q_{0E}^2}{2} \right] + D_E q_{0E} \frac{\mathbf{p} \cdot \mathbf{q}}{m} \right] \\
T_2 &= \tilde{\theta}_E(q_E) \tilde{\theta}_E(k_E) \tilde{\theta}_E(-q_E - k_E) A_{E-} A_{E+} \frac{-(\mathbf{k} \cdot \mathbf{q} + \mathbf{k}^2)}{8m} \left[[B_E + C_E] q_{0E} + D_E \frac{\mathbf{p} \cdot \mathbf{q}}{m} \right] \\
T_3 &= \tilde{\theta}_E(k_E) \tilde{\theta}_E(r_E) \tilde{\theta}_E(q_E - k_E) \tilde{\theta}_E(-q_E - r_E) A_{E-} A_{E+} \frac{-\mathbf{k}^2 + \mathbf{k} \cdot \mathbf{q}}{32m^2} (\mathbf{r}^2 + \mathbf{r} \cdot \mathbf{q}) [B_E + C_E]
\end{aligned}$$

with

$$\begin{aligned}
A_E(p_E) &= \frac{-1}{p_{0E}^2 + E(\mathbf{p})^2}, \quad A_{E+} = A_E(p_E + \frac{q_E}{2}), \quad A_{E-} = A_E(p_E - \frac{q_E}{2}) \\
B_E(p_{0E}, q_{0E}, \mathbf{p}, \mathbf{q}) &= |\Delta|^2 \\
C_E(p_{0E}, q_{0E}, \mathbf{p}, \mathbf{q}) &= -\epsilon(\mathbf{p} - \frac{\mathbf{q}}{2}) \epsilon(\mathbf{p} + \frac{\mathbf{q}}{2}) + (p_{0E}^2 - (\frac{q_{0E}}{2})^2) \\
D_E(p_{0E}, q_{0E}, \mathbf{p}, \mathbf{q}) &= \epsilon(\mathbf{p} + \frac{\mathbf{q}}{2}) (p_{0E} - \frac{q_{0E}}{2}) + \epsilon(\mathbf{p} - \frac{\mathbf{q}}{2}) (p_{0E} + \frac{q_{0E}}{2}).
\end{aligned}$$

We note that the integrals over p_{0E} all have the form

$$\tilde{g}_E = \int_{-\infty}^{\infty} \frac{dp_{0E}}{2\pi} A_{E-} A_{E+} g_E(p_{0E}, q_{0E}, \mathbf{p}, \mathbf{q}) \quad g = B, C, D. \quad (20)$$

We now proceed to the second step of the finite temperature formalism which is to discretise the momentum variables as follows

$$\begin{aligned}
p_{0E} \rightarrow p_{0E}^n &= \frac{(2n+1)\pi}{\beta}, \quad q_{0E}/2 \rightarrow q_{0E}^m/2 = \frac{(2m\pi)}{\beta} \\
k_{0E} \rightarrow k_{0E}^l &= \frac{(2l\pi)}{\beta}, \quad r_{0E} \rightarrow r_{0E}^s = \frac{(2s\pi)}{\beta}
\end{aligned} \quad (21)$$

and then calculate the sums over n which are contained in (20) by replacing them with integrals over z according to (13). The poles of $\tan(\frac{\beta z}{2})$ are on the real axis and the sum of their residues is equal to the finite temperature sum which is to be calculated. We note that only A, B, C, D depend on the variable z and B, C, D have no poles. This means that we need to evaluate only the following integrals

$$\tilde{g}_E = \int_C dz A_E(z + \frac{q_{0E}^m}{2}, \mathbf{p} + \frac{\mathbf{q}}{2}) A_E(z - \frac{q_{0E}^m}{2}, \mathbf{p} - \frac{\mathbf{q}}{2}) g_E(z, q_{0E}^m, \mathbf{p}, \mathbf{q}) \tan(\frac{\beta z}{2}) \quad (22)$$

where the function g_E has no poles (analytical everywhere in the z -plane) and the A -functions have the poles

$$\frac{q_{0E}^m}{2} + iE_-, \frac{q_{0E}^m}{2} - iE_-, -\frac{q_{0E}^m}{2} + iE_+, -\frac{q_{0E}^m}{2} - iE_+$$

where we have introduced the usual notation

$$E_+ = E(\mathbf{p} + \frac{\mathbf{q}}{2}) \quad E_- = E(\mathbf{p} - \frac{\mathbf{q}}{2}).$$

We follow the same procedure as for the first trace. We note from (21) that $\exp(i\beta \frac{q_{0E}^m}{2}) = 1$ and then, following [7, 9, 14], we Wick rotate the above expression back to real continuous variables q_0 via $q_{0E}^m \rightarrow -iq_0$. The result can be written as $\tilde{g} = \tilde{g}_R + \tilde{g}_L$, where the regular part $\tilde{g}_R$ has a well-defined expansion about the origin $(q_0, \mathbf{q}) = (0, \mathbf{0})$, but the Landau part $\tilde{g}_L$ does not, as was first pointed out in [8]. For $g = B, C, D$ we have

$$\begin{aligned} \tilde{B}_R &= \frac{|\Delta|^2}{4E_+E_-} [1 - n(E_+) - n(E_-)] \left(\frac{1}{E_+ + E_- - q_0} + \frac{1}{E_+ + E_- + q_0} \right) \\ \tilde{B}_L &= \frac{|\Delta|^2}{2E_+E_-} \frac{n(E_+) - n(E_-)}{E_+ - E_- + q_0} \\ \tilde{C}_R &= F_1 [1 - n(E_+) - n(E_-)] \left(\frac{1}{E_+ + E_- - q_0} + \frac{1}{E_+ + E_- + q_0} \right) \\ \tilde{C}_L &= 2F_2 \frac{n(E_-) - n(E_+)}{E_+ - E_- + q_0} \\ \tilde{D}_R &= G_1 [1 - n(E_+) - n(E_-)] \left(\frac{1}{E_+ + E_- - q_0} - \frac{1}{E_+ + E_- + q_0} \right) \\ \tilde{D}_L &= 2G_2 \frac{n(E_+) - n(E_-)}{E_+ - E_- + q_0} \end{aligned}$$

with

$$F_1 = \frac{1}{4} \left[1 - \frac{\epsilon_+ \epsilon_-}{E_+ E_-} \right], \quad F_2 = \frac{1}{4} \left[1 + \frac{\epsilon_+ \epsilon_-}{E_+ E_-} \right]$$

$$G_1 = \frac{i}{2} \frac{\epsilon_+}{E_+}, \quad G_2 = \frac{i}{4} \left[\frac{\epsilon_+}{E_+} + \frac{\epsilon_-}{E_-} \right].$$

After the analytic continuation, wherever we write q_0 in the denominators of B, C, D , we imply that there is an infinitesimal imaginary part, $i\delta$, $\delta \rightarrow 0^+$ added to q_0 , corresponding to retarded

boundary conditions [7, 9, 15]. We complete the Wick rotation back to continuous variables via $k_{0E}^m \rightarrow -ik_0$, $r_{0E}^m \rightarrow -ir_0$, and the second trace finally takes the form

$$\begin{aligned}
\text{Tr}[\hat{\Sigma}\hat{G}\hat{\Sigma}\hat{G}] &= i \int \frac{d^4q}{(2\pi)^4} \tilde{\theta}(q) \tilde{\theta}(-q) \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[\tilde{B} \left[\frac{(\mathbf{p} \cdot \mathbf{q})^2}{2m^2} - \frac{q_0^2}{2} \right] \right. \\
&\quad \left. + \tilde{C} \left[-\frac{(\mathbf{p} \cdot \mathbf{q})^2}{2m^2} - \frac{q_0^2}{2} \right] + \tilde{D} (-i)q_0 \frac{\mathbf{p} \cdot \mathbf{q}}{m} \right] \\
&+ 2i \int \frac{d^4q}{(2\pi)^4} \frac{d^4k}{(2\pi)^4} \tilde{\theta}(q) \tilde{\theta}(k) \tilde{\theta}(-q-k) \frac{-(\mathbf{k} \cdot \mathbf{q} + \mathbf{k}^2)}{8m} \\
&\quad \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[[\tilde{B} + \tilde{C}](-i)q_0 + \tilde{D} \frac{\mathbf{p} \cdot \mathbf{q}}{m} \right] \\
&+ i \int \frac{d^4q}{(2\pi)^4} \frac{d^4k}{(2\pi)^4} \frac{d^4r}{(2\pi)^4} \tilde{\theta}(k) \tilde{\theta}(r) \tilde{\theta}(q-k) \tilde{\theta}(-q-r) \\
&\quad \frac{(-\mathbf{k}^2 + \mathbf{k} \cdot \mathbf{q})(\mathbf{r}^2 + \mathbf{r} \cdot \mathbf{q})}{32m^2} \int \frac{d^3\mathbf{p}}{(2\pi)^3} [\tilde{B} + \tilde{C}]. \tag{23}
\end{aligned}$$

5 The effective Lagrangian at $T \neq 0$, neglecting the Landau terms

The contribution of the first Trace to the effective action is given in (16). When the Landau terms are neglected, only the functions $\tilde{B}_R$, $\tilde{C}_R$ and $\tilde{D}_R$ appear in the expression (23) for the second Trace and they can all be Taylor-expanded about $q = 0$ in powers of $q_0/|\Delta|$ and $v_F|\mathbf{q}|/|\Delta|$, so long as these quantities are small. For $\tilde{B}_R$ and $\tilde{C}_R$ the expansion takes the form

$$\tilde{B}_R(\mathbf{p}, q) = \frac{|\Delta|^2}{4E^3}(1 - 2n(E)) + O(\mathbf{q}^2, (\mathbf{q} \cdot \mathbf{p})^2, q_0^2) \tag{24}$$

$$\tilde{C}_R(\mathbf{p}, q) = \frac{|\Delta|^2}{4E^3}(1 - 2n(E)) + O(\mathbf{q}^2, (\mathbf{q} \cdot \mathbf{p})^2, q_0^2). \tag{25}$$

$\tilde{D}_R(q)$, on the other hand, vanishes as $q = 0$, and we have

$$\tilde{D}_R(\mathbf{p}, q) = O(q_0(\mathbf{q} \cdot \mathbf{p})). \tag{26}$$

When (24) - (26) are inserted in (23) and the appropriate Fourier transforms performed, we

obtain a local effective Lagrangian for θ , involving powers and products of the derivatives θ and $\nabla\theta$.

We must now consider how this derivative expansion is to be ordered. Due to the preferred frame supplied by the heat bath, we do not expect Galilean invariance to be preserved in the effective Lagrangian at $T \neq 0$. Nevertheless, we expect Galilean invariance to remain approximately valid for $T \leq T_c$. Accordingly we shall regard terms of the form “ $\dot{\theta}$ ” and “ $(\nabla\theta)^2/m$ ” as being of the same order - or equivalently, terms of the form “ $q_0\theta$ ” and “ $\mathbf{q}^2\theta^2$ ” as the same order. It then follows that the leading order terms in the expansion of (23) are given by retaining only the *zeroth* order contributions to $\tilde{B}_R, \tilde{C}_R$ and $\tilde{D}_R$ from (24)-(26), thanks to the explicit factors involving q_0 and $\mathbf{q}$ already appearing in (23). This is an important point.

In evaluating these leading order terms, it is convenient to define

$$N(T) = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{|\Delta|^2}{2E^3} (1 - 2n(E)), \quad (27)$$

and to make use of the approximation [9]

$$N(T) \approx \frac{1}{2} N(0) R(T) \quad (28)$$

where

$$N(0) = \frac{mp_F}{2\pi^2} = \frac{3\rho_0}{2mv_F^2} \quad (29)$$

is the density of states at the Fermi energy and

$$R(T) = \int dy \frac{1 - 2n(y)}{(1 + y^2)^{\frac{3}{2}}} \quad (30)$$

where $y = \epsilon / |\Delta|$. The leading contribution to the first term in (23) is then

$$-i\frac{1}{2}N(T) \int \frac{d^4q}{(2\pi)^4} \tilde{\theta}(q)\tilde{\theta}(-q)q_0^2 = -\frac{i}{2}N(T) \int d^4x \dot{\theta}^2(x). \quad (31)$$

Evaluating the other terms in the same way, and including the expression (16) for the first Trace, we obtain the effective Lagrangian (neglecting Landau terms)

$$\mathcal{L}_R(\theta) = -M(T) \left[\dot{\theta} + \frac{(\nabla\theta)^2}{4m} \right] + \frac{N(T)}{4} \left[\dot{\theta} + \frac{(\nabla\theta)^2}{4m} \right]^2. \quad (32)$$

Remarkably, Galilean invariance is still preserved by (32). The only difference between (32) and $\mathcal{L}_{eff}(\theta)$ of [1, 4] is the appearance of the T -dependent coefficients $M(T)$ and $N(T)$. In fact, as $T \rightarrow 0$ we note that $M(T) \rightarrow \rho_0/2$, while $N(T) \rightarrow N(0)$, so the $T = 0$ result of [1, 4] is easily regained.

The Galilean invariance of (32) guarantees that it can be rewritten as a time-dependent non-linear Schrödinger Lagrangian (TDNLSL), as explained in [4]. The results of [4] can be immediately taken over to $T \neq 0$ (always with the neglect of the Landau terms) by making the replacements $\rho_0/2 \rightarrow M(T)$, $N(0) \rightarrow N(T)$. The result is merely that the “potential” term in the equivalent TDNLSL becomes

$$V = \frac{[\rho - 2M(T)]^2}{2N(T)}. \quad (33)$$

Thus, always with neglect of Landau terms, we have shown that the effective theory for our neutral fermionic fluid is a TDNLSL, just as it is for a bosonic fluid. The inclusion of electromagnetism by means of the “minimal coupling” prescription is straightforward.

We now turn to the Landau terms which, as in [10], are non-local and cannot be Taylor expanded about $q = 0$.

6 The Landau terms

The Landau terms $\tilde{B}_L$, $\tilde{C}_L$ and $\tilde{D}_L$ cannot be treated in the same way as the regular terms, because of their non-analyticity around the origin. However, since we are interested in the low frequency, long wavelength regime, we can follow [8, 14] and write

$$E_+ - E_- = \frac{\epsilon}{E} \frac{\mathbf{p} \cdot \mathbf{q}}{m}. \quad (34)$$

We now expand the numerators of these quantities in powers of momentum $\mathbf{q}$ and keep the leading order terms which are of first order in $\mathbf{q}$. We find

$$\tilde{B}_L \approx \left[\frac{\epsilon}{E} \frac{\mathbf{p} \cdot \mathbf{q}}{m} + q_0 + i\delta \right]^{-1} \frac{|\Delta|^2}{2E^2} \frac{dn}{dE} \frac{\epsilon}{E} \frac{\mathbf{p} \cdot \mathbf{q}}{m}$$

$$\begin{aligned}
\tilde{C}_L &\approx - \left[\frac{\epsilon}{E} \frac{\mathbf{p} \cdot \mathbf{q}}{m} + q_0 + i\delta \right]^{-1} \frac{2\epsilon^2 + |\Delta|^2}{2E^2} \frac{dn}{dE} \frac{\epsilon}{E} \frac{\mathbf{p} \cdot \mathbf{q}}{m} \\
\tilde{D}_L &\approx \left[\frac{\epsilon}{E} \frac{\mathbf{p} \cdot \mathbf{q}}{m} + q_0 + i\delta \right]^{-1} \frac{dn}{dE} \frac{i\epsilon^2}{E^2} \frac{\mathbf{p} \cdot \mathbf{q}}{m}.
\end{aligned} \tag{35}$$

If we look at (23) and recall that $\tilde{g} = \tilde{g}_R + \tilde{g}_L$ for $\tilde{g} = \tilde{B}, \tilde{C}, \tilde{D}$ we realize that the Landau terms appear in the second Trace in the following integrals

$$\begin{aligned}
I_1 &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[\tilde{B}_L \left[\frac{(\mathbf{p} \cdot \mathbf{q})^2}{2m^2} - \frac{q_0^2}{2} \right] + \tilde{C}_L \left[-\frac{(\mathbf{p} \cdot \mathbf{q})^2}{2m^2} - \frac{q_0^2}{2} \right] + \tilde{D}_L (-i)q_0 \frac{\mathbf{p} \cdot \mathbf{q}}{m} \right] \\
I_2 &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[[\tilde{B}_L + \tilde{C}_L] (-i)q_0 + \tilde{D}_L \frac{\mathbf{p} \cdot \mathbf{q}}{m} \right] \\
I_3 &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} [\tilde{B}_L + \tilde{C}_L].
\end{aligned} \tag{36}$$

Substituting into I_1 the expressions (35) for $\tilde{B}_L, \tilde{C}_L, \tilde{D}_L$ we obtain

$$I_1 = \int \frac{|\mathbf{p}|^2 d|\mathbf{p}|}{4\pi^2} \frac{dn}{dE} \left[\frac{|\mathbf{p}|^2 |\mathbf{q}|^2}{3m^2} + \frac{|\Delta|^4}{\epsilon^2 E^2} \left(1 + \frac{a}{2} I_s \right) q_0^2 \right], \tag{37}$$

where

$$I_s = \int_{-1}^1 \frac{ds}{s - a - i\delta \operatorname{sgn}(\epsilon)}, \quad a = \frac{q_0 E m}{\epsilon |\mathbf{p}| |\mathbf{q}|}.$$

The integral I_s has a pole at a , if $-1 < a < 1$. The sign of ϵ appears in the denominator of I_s , because the infinitesimal imaginary part $i\delta$ is multiplied by a whose sign depends only on the sign of ϵ , the rest of the parameters in a being always positive. Performing the s -integration yields

$$I_s = \ln \left| \frac{1-a}{1+a} \right| + i\pi \operatorname{sgn}(\epsilon) \theta(1-|a|). \tag{38}$$

Similar expressions can be found for I_2 and I_3 . Applying now approximation (28) we find

$$\begin{aligned}
I_1 &= \left(\frac{3\rho_0}{4m} \right) \left(\frac{1}{3} J |\mathbf{q}|^2 - (H_r^{(1)}(c) + iH_i^{(1)}(c)) q_0^2 / v_F^2 \right) \\
I_2 &= i \left(\frac{3\rho_0}{2m v_F^2} \right) q_0 (H_r^{(2)}(c) + iH_i^{(2)}(c)) \\
I_3 &= \left(\frac{3\rho_0}{2m v_F^2} \right) (H_r^{(3)}(c) + iH_i^{(3)}(c))
\end{aligned} \tag{39}$$

where

$$\begin{aligned}
J &= |\Delta| \int dy \frac{dn}{dE} \\
H_r^{(n)}(c) &= -|\Delta| \int dy \frac{dn}{dE} \frac{y^{2(n-2)}}{(y^2+1)} \left[1 - \frac{c}{2} \frac{\sqrt{y^2+1}}{y} \ln \left| \frac{y+c\sqrt{y^2+1}}{y-c\sqrt{y^2+1}} \right| \right] \\
H_i^{(n)}(c) &= -\frac{|\Delta|\pi c}{2} \int dy \frac{dn}{dE} \frac{y^{2(n-2)}}{|y|\sqrt{1+y^2}} \theta\left(\frac{|y|}{\sqrt{1+y^2}} - |c|\right) \\
c &= \frac{1}{v_F} \frac{q_0}{|\mathbf{q}|}.
\end{aligned} \tag{40}$$

We observe that, for $|c| \geq 1$, $H_i^{(n)} = 0$, so that the cut in the $H^{(n)}$ functions is over the region $|c| \leq 1$. It is noteworthy that the function $H^{(1)}$, which multiplies the term in q_0^2 , is, apart from a factor of 4 and slight changes in notation, exactly the same as the corresponding function H of [10].

Substituting the expressions which we have found for I_1, I_2, I_3 back into the second trace, making approximation (17), and Fourier transforming the $\tilde{\theta}$ functions wherever this is possible, we can bring the effective action to the following form

$$\begin{aligned}
S_{eff} &= \int d^4x \left(-\frac{3\rho_0}{8m} \right) \left[\frac{4m}{3} \left(\dot{\theta} + [1+J] \frac{(\nabla\theta)^2}{4m} \right) - \frac{R}{2v_F^2} \left(\dot{\theta} + \frac{(\nabla\theta)^2}{4m} \right)^2 \right. \\
&\quad - \int \frac{d^4q}{(2\pi)^4} \tilde{\theta}(q) \tilde{\theta}(-q) \frac{q_0^2}{v_F^2} (H_r^{(1)}(c) + iH_i^{(1)}(c)) \\
&\quad - i \int \frac{d^4q}{(2\pi)^4} \frac{d^4k}{(2\pi)^4} \tilde{\theta}(q) \tilde{\theta}(k) \tilde{\theta}(-q-k) 2q_0 \frac{(\mathbf{k}\cdot\mathbf{q} + \mathbf{k}^2)}{4m} \frac{1}{v_F^2} (H_r^{(2)}(c) + iH_i^{(2)}(c)) \\
&\quad + \int \frac{d^4q}{(2\pi)^4} \frac{d^4k}{(2\pi)^4} \frac{d^4r}{(2\pi)^4} \tilde{\theta}(k) \tilde{\theta}(r) \tilde{\theta}(q-k) \tilde{\theta}(-q-r) \\
&\quad \left. \frac{(-\mathbf{k}^2 + \mathbf{k}\cdot\mathbf{q})}{4m} \frac{(\mathbf{r}^2 + \mathbf{r}\cdot\mathbf{q})}{4m} \frac{1}{v_F^2} (H_r^{(3)}(c) + iH_i^{(3)}(c)) \right].
\end{aligned} \tag{41}$$

Consider now the effect of the Landau terms J and $H^{(n)}$ in (41). The simplest of these is J , which is a (temperature-dependent) constant. Thus its contribution is still local but its presence does break Galilean invariance, since it disrupts the invariant combination “ $\dot{\theta} + (\nabla\theta)^2/4m$ ” in the first term in square brackets in (41). Actually, as far as this single term is concerned, Galilean invariance can be formally restored by modifying the first $\dot{\theta}$ term so as to read $[1+J]\dot{\theta}$; this does not affect the equations of motion.

The effect of the $H^{(n)}$ functions is - depending on their numerical magnitudes - potentially much more drastic. Most importantly, these functions cannot be expanded about $q = 0$, since they depend explicitly on the ratio $c = q_0/v_F|\mathbf{q}|$. Thus the terms involving the $H^{(n)}$'s in (41) cannot be represented as any Lagrangian local in coordinate space. *A fortiori* then, these terms also break Galilean invariance. They also contain imaginary parts, of course, corresponding to the physical “ Cerenkov ” radiation process in the heat bath [8, 10]. In the next section we discuss how the equation of motion for θ , which follows from (41), may be approximated by a more manageable form, which may be given a physical interpretation.

7 Approximate equation of motion for θ and its interpretation

7.1 The θ -propagator

Let us begin by considering the terms in (41) which are quadratic in $\tilde{\theta}$, from which we can read off the momentum-space propagator for θ . The first point to note is that, since $H_i^{(1)}$ is an odd function of q_0 , it contributes nothing to the action S_{eff} . To see the effect of damping we need to consider the equation of motion for $\tilde{\theta}$. This has the form

$$\tilde{\theta} = S_G(q_0, |\mathbf{q}|) \left[\int \tilde{\theta}^2 H_{r,i}^{(2)}(c), \int \int \tilde{\theta}^3 H_{r,i}^{(3)}(c) \right] \quad (42)$$

where we have written the non-linear terms on the right hand side in symbolic form and where the θ -propagator is given by

$$S_G(q_0, |\mathbf{q}|) = \left\{ \mathbf{q}^2 \left[\frac{c^2 R}{2} - \frac{1+J}{3} + c^2 \left(H_r^{(1)}(c) + i H_i^{(1)}(c) \right) \right] \right\}^{-1} \quad (43)$$

At $T = 0$ we have $R = 2$ and $J = H_r^{(1)} = H_i^{(1)} = 0$, so that S_G has poles at $c = \pm \frac{1}{\sqrt{3}}$, as expected for the Bogoliubov-Anderson mode [16, 17].

Consider now what happens as T moves away from zero. $\frac{R}{2}$ and J deviate slowly from 1 and 0 respectively (see fig.1a, fig.1b) while the quantities $c^2 H_r^{(1)}$ and $c^2 H_i^{(1)}$ are very small compared to the main term $-\frac{1}{3}$ (see fig.2a, fig.2b). This suggests that it should be a good approximation

to replace the complicated c -dependence of the denominator in (43) by a quadratic function, which amounts to replacing the cut structure of (43) by two poles - the same two that were present at $T = 0$ but now at slightly different positions. In particular, the poles will now be shifted off the real axis, due to the presence of the $H_i^{(1)}$ function. It is important to note that $H_r^{(1)}$ is an even function of real c while $H_i^{(1)}$ is an odd function of real c . It is then clear that the poles of (43) are approximately at $c = \pm c_r - i\gamma$, where $c_r = \pm \frac{1}{\sqrt{3}}$ and $\gamma \ll c_r$; both of the approximate poles are in the lower half plane, as expected, since we are considering a retarded propagator, which is required for the study of damping.

To find expressions for c_r and γ we write $H_i^{(1)}(c) = c \tilde{H}_i^{(1)}(c^2)$, so that, after the approximation which follows, we do not lose the property that $H_i^{(1)}$ is an odd function of c . Now, since the functions $H_r^{(1)}$ and $H_i^{(1)}$ are small compared to the regular term, we can make an approximation similar to the one which leads to the well-known Breit-Wigner formula; we expand the quantities $c^2 H_r^{(1)}(c^2)$ and $c^2 \tilde{H}_i^{(1)}(c^2)$ which appear in (43) about $c^2 = \frac{1}{3}$ keeping only the lowest order term. Thus we arrive at

$$S_G(q_0, |\mathbf{q}|) \approx \left\{ \mathbf{q}^2 \frac{R}{2} \left[c^2 - \frac{2(1+J)}{3R} + \frac{2\bar{H}_r^{(1)}}{3R} + i \frac{2c \bar{H}_i^{(1)}}{\sqrt{3}R} \right] \right\}^{-1} \quad (44)$$

where the bar denotes quantities evaluated at $c = \frac{1}{\sqrt{3}}$. (44) exhibits the desired quadratic behaviour and has poles at $c = \pm c_r - i\gamma$, where

$$c_r = \sqrt{\frac{2(1+J)}{3R} - \frac{2\bar{H}_r^{(1)}}{3R}} \quad \gamma = \frac{\bar{H}_i^{(1)}}{\sqrt{3}R}$$

and γ^2 has been neglected. c_r and γ , as functions of temperature, are plotted in fig.3a and fig.3b respectively. They both have the expected behaviour, which reassures us about the validity of our approximation. In terms of c_r and γ we may recast (44) as follows

$$S_G(q_0, |\mathbf{q}|) \approx \left\{ \frac{R}{2v_F^2} \left[q_0^2 - c_r^2 v_F^2 \mathbf{q}^2 + 2i \gamma v_F q_0 |\mathbf{q}| \right] \right\}^{-1} \quad (45)$$

or equivalently as

$$S_G \approx \frac{v_F}{c_r R |\mathbf{q}|} \left\{ \frac{1}{q_0 - c_r v_F |\mathbf{q}| + i\gamma v_F |\mathbf{q}|} - \frac{1}{q_0 + c_r v_F |\mathbf{q}| + i\gamma v_F |\mathbf{q}|} \right\} \quad (46)$$

which manifestly has the correct form for a bosonic propagator with damping, the line-width having an explicit $|\mathbf{q}|$ -dependence.

At this stage we have to point out that the alternative approximate form for S_G given in [10] is incorrect, since it fails to exhibit the important property (traceable to $H_i^{(1)}$) that the imaginary part should be an odd function of q_0 . The pole of S_G at $c = -c_r - i\gamma$ was overlooked in [10] and instead a pole at $c = -c_r + i\gamma$ was used, which is actually on a Riemann sheet distant from the physical boundary. This mistake had the effect of replacing the $2i \gamma v_F q_0 |\mathbf{q}|$ term in (45) with something proportional to $\mathbf{q}^2$. Such a form then does, of course, have a simple local Fourier transform. In contrast (45) does not but, as noted in the start of the section, the odd- q_0 piece will integrate to zero in S_{eff} . Damping can only be manifested in the equation of motion and in this case, when included, it will lead to a non-local term in the effective wave-operator (the transform of S_G^{-1}). Nevertheless, we do have a simple and reliable expression (45) for S_G in momentum space, which shows a particular form of damping and a predicted temperature-dependence for the line-shape. As noted in the Introduction, the non-local form of S_G^{-1} appears to present a fundamental difficulty for the inclusion of electromagnetism.

We now turn to the treatment of the other terms on the right hand side of (42).

7.2 The remaining Landau terms involving $H^{(2)}$ and $H^{(3)}$.

The fact that S_G is so sharply peaked around $q_0 \approx \pm \frac{1}{\sqrt{3}} v_F |\mathbf{q}|$ suggests that, as far as the dynamics of θ described by (42) are concerned, it should be a good approximation to evaluate the c -even functions $H_r^{(2)}$ and $H_r^{(3)}$ at $c^2 = \frac{1}{3}$. The corresponding imaginary parts are very small, c -odd functions, as $H_i^{(1)}$ was, and we shall from now on simply ignore them. In this case all the terms in (41) become local and the Fourier transforms to x -space can all be done, leading to the effective Lagrangian

$$\begin{aligned} \mathcal{L}_{R+\bar{L}_i} = & -M'(T) \left[\dot{\theta} + \frac{(\nabla\theta)^2}{4m} \right] + \frac{N(T)}{4} \left[\dot{\theta} + \frac{(\nabla\theta)^2}{4m} \right]^2 \\ & + \frac{3\rho_0}{8mv_F^2} \left(\dot{\theta}^2 \bar{H}_r^{(1)} + 2\dot{\theta} \frac{(\nabla\theta)^2}{4m} \bar{H}_r^{(2)} + \frac{[(\nabla\theta)^2]^2}{(4m)^2} \bar{H}_r^{(3)} \right), \end{aligned} \quad (47)$$

where

$$M'(T) = \frac{\rho_0}{2} (1 + J)$$

is (17) as modified by the contribution from the Landau term, $\bar{H}_r^{(j)} \equiv H_r^{(j)}(c^2 = \frac{1}{3})$ ($j = 1, 2, 3$) and we have modified the $\dot{\theta}$ term as suggested after (41). We note that the first part of (47),

not involving the $\bar{H}_r^{(j)}$ ($j = 1, 2, 3$) can be written as a TDNLSL just as in section 5, with a potential

$$V = \frac{(\rho - 2M')^2}{2N}.$$

Choosing a convenient normalisation, the equation of motion for θ is then

$$\begin{aligned} & \frac{\partial}{\partial t} \left[\rho - \frac{3\rho_0}{2mv_F^2} (\dot{\theta} \bar{H}_r^{(1)} + \frac{(\nabla\theta)^2}{4m} \bar{H}_r^{(2)}) \right] \\ & + \nabla \cdot \left\{ \frac{\nabla\theta}{2m} \left[\rho - \frac{3\rho_0}{2mv_F^2} \left(\dot{\theta} \bar{H}_r^{(2)} + \frac{(\nabla\theta)^2}{4m} \bar{H}_r^{(3)} \right) \right] \right\} = 0 \end{aligned} \quad (48)$$

where

$$\rho = 2M' - N \left[\dot{\theta} + \frac{(\nabla\theta)^2}{4m} \right]. \quad (49)$$

It is tempting to interpret the quantity

$$\rho_n = -\frac{3\rho_0}{2mv_F^2} \left(\dot{\theta} \bar{H}_r^{(1)} + \frac{(\nabla\theta)^2}{4m} \bar{H}_r^{(2)} \right) \quad (50)$$

in (48) as an additional “density”, not described by the NLSL. On this interpretation, we would expect the corresponding additional term inside the divergence in (48) to represent the associated “current”, which would require

$$\rho_n \mathbf{v}_n = -\frac{3\rho_0}{2mv_F^2} \left(\dot{\theta} \bar{H}_r^{(2)} + \frac{(\nabla\theta)^2}{4m} \bar{H}_r^{(3)} \right) \frac{\nabla\theta}{2m}. \quad (51)$$

Equations (50) and (51) are compatible, if

$$\mathbf{v}_n = \frac{\nabla\theta}{2m^*} \quad (52)$$

$$m^* = (\bar{H}_r^{(1)} / \bar{H}_r^{(2)}) m \quad (53)$$

$$\bar{H}_r^{(3)} \bar{H}_r^{(1)} = (\bar{H}_r^{(2)})^2. \quad (54)$$

In this case, (48) becomes the continuity equation of the two-fluid model

$$\frac{\partial}{\partial t} (\rho + \rho_n) + \nabla \cdot (\rho \mathbf{v} + \rho_n \mathbf{v}_n) = 0 \quad (55)$$

where ρ is the superfluid density and the superfluid velocity is $\mathbf{v} = \nabla\theta/2m$ as usual.

The crucial requirement is (54). Fig.4a shows the ratio $\bar{H}_r^{(3)} \bar{H}_r^{(1)} / (\bar{H}_r^{(2)})^2$ over the range of b we are considering, and it can be seen that (54) is quite well satisfied. Fig.4b shows the ratio

$\bar{H}_r^{(1)}/\bar{H}_r^{(2)}$ as a function of b , from which we see that m^* in (53) varies quite rapidly with T at low T , and then approaches the constant value $5m$.

It is easy to show that there is a TDNLSL for the normal fluid from which the continuity equation for the normal fluid can be derived as an equation of motion. One can then use the “minimal coupling” prescription to take electromagnetism into account.

In summary, we have argued that the effect of the Landau terms $H^{(n)}$ can be approximately incorporated by treating them as a constant, evaluating them at $c^2 = 1/3$. Considering the resulting effective Lagrangian, it seems to us to describe a two-fluid model in which the superfluid component could be described by a TDNLSL with a temperature dependent potential, and the normal component by another TDNLSL with an appropriately chosen potential and a definite velocity predicted by (52).

Acknowledgements

We wish to thank Cesar Fosco, Marcelo Hott and Alexei Tsvelik for valuable discussions. I.J.R.A. gratefully acknowledges support from the Antorchas Foundation (Argentina) and the British Council, which enabled him to visit the Instituto Balseiro, Bariloche, where part of this work was completed. D.J.L. and G.M. gratefully acknowledge financial support from P.P.A.R.C. (UK).

References

- [1] A. M. J. Schakel, Mod.Phys.Lett. **B4**, 927 (1990).
- [2] C. M. Fraser, Z. Phys. **C28**, 101 (1985).
- [3] M. Greiter, F. Wilczek, and E. Witten, Mod. Phys. Lett. **B3**, 903 (1989).
- [4] I. J. R. Aitchison, P. Ao, D. J. Thouless, and X. M. Zhu, Phys.Rev. **B51**, 6531 (1995).
- [5] M. Stone, Int.J.Mod.Phys **B9**, 1359 (1995).
- [6] S. D. Palo, C. Castellani, C. D. Castro, and B. K. Chakraverty, cond.-mat/9901305 (1999).
- [7] J. R. Schrieffer, *Theory of Superconductivity* (Addison-Wesley, New York, 1988).
- [8] E. Abrahams and T. Tsuneto, Phys.Rev. **152**, 416 (1966).
- [9] H. T. C. Stoof, Phys.Rev. **B47**, 7979 (1992).
- [10] I. J. R. Aitchison and D. J. Lee, Phys.Rev. **B56**, 8303 (1997).
- [11] C. G. van Weert, *Thermal Field Theories* (Elsevier Science Publisers, B.V., 1991), p. 17.
- [12] M. Houbiers and H. T. C. Stoof, cond-mat/9804241 (1998).
- [13] J. Bardeen, L. N. Cooper, and J. R. Schrieffer, Phys. Rev. **108**, 1175 (1957).
- [14] I. J. R. Aitchison and J. A. Zuk, Ann. Phys. **242**, 77 (1995).
- [15] H. A. Weldon, Phys. Rev. **D47**, 594 (1993).
- [16] P. Anderson, Phys.Rev. **112**, 1900 (1958).
- [17] N. N. Bogoliubov, V. V. Tolmachev, and D. V. Shirkov, *A New Method in the Theory of Superconductivity* (Consultants Bureau, NY, 1959).




Figure 1: (a): $\frac{R(T)}{2}$ is very close to 1 for any temperature in the region which is considered here, $b = \frac{T}{T_c} \in [0, 0.6]$. (b): $|J(T)|$ is much smaller than 1 for any temperature in the same region.




Figure 2: (a): $c^2|H_r^{(1)}|$ is negligible compared to $\frac{1}{3} \approx 0.333$, for any temperature $b \in [0, 0.6]$ and $|c| \leq 1$. For $|c| \geq 1$, it can be shown that the c -even function $c^2|H_r^{(1)}|$ is always much smaller than $\frac{1}{3}$.




Figure 2: (b): $c^2 |H_i^{(1)}|$ is negligible compared to $\frac{1}{3} \approx 0.333$, for any temperature $b \in [0, 0.6]$ and $|c| \leq 1$. For $|c| \geq 1$, $H_i^{(1)} = 0$.




Figure 3: (a): c_r is very close to the original location of the absolute value of the poles $\frac{1}{\sqrt{3}} \approx 0.577$, for any temperature in the considered region. (b): The damping γ is negligible compared to the real part of the pole, for any temperature in the considered region.




Figure 4: (a): As the temperature rises $\sqrt{\bar{H}_r^{(1)} \bar{H}_r^{(3)}}/|\bar{H}_r^{(2)}|$ approaches 1. (b): As the temperature rises $\bar{H}_r^{(1)}/\bar{H}_r^{(2)} = m^*/m$ approaches 5.